

Weak type estimates of Marcinkiewicz integrals on the weighted Hardy and Herz-type Hardy spaces

Hua Wang*

School of Mathematical Sciences, Peking University, Beijing 100871, China

Abstract

The Marcinkiewicz integral is essentially a Littlewood-Paley g -function, which plays an important role in harmonic analysis. In this article, by using the atomic decomposition theory of weighted Hardy spaces and homogeneous weighted Herz-type Hardy spaces, we will obtain some weighted weak type estimates for Marcinkiewicz integrals on these spaces.

MSC(2000): 42B25; 42B30

Keywords: Marcinkiewicz integrals; weighted Hardy spaces; weighted Herz-type Hardy spaces; A_p weights; atomic decomposition

1 Introduction

Suppose that S^{n-1} is the unit sphere in $\mathbb{R}^n$ ($n \geq 2$) equipped with the normalized Lebesgue measure $d\sigma$. Let $\Omega \in L^1(S^{n-1})$ be homogeneous of degree zero and satisfy the cancellation condition

$$\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0,$$

where $x' = x/|x|$ for any $x \neq 0$. Then the Marcinkiewicz integral of higher dimension μ_Ω is defined by

$$\mu_\Omega(f)(x) = \left(\int_0^\infty |F_{\Omega,t}(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,t}(x) = \int_{|x-y| \leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy.$$

*E-mail address: wanghua@pku.edu.cn.

This operator μ_Ω was first introduced by Stein in [14]. He proved that if $\Omega \in Lip_\alpha(S^{n-1})$ ($0 < \alpha \leq 1$), then μ_Ω is of type (p, p) for $1 < p \leq 2$ and of weak type $(1, 1)$. It is well known that the Littlewood-Paley g -function is a very important tool in harmonic analysis and the Marcinkiewicz integral is essentially a Littlewood-Paley g -function. Therefore, many authors have been interested in studying the boundedness properties of μ_Ω on various function spaces, we refer the readers to see [1, 2, 3, 7, 9, 16] for its developments and applications.

In 1990, Torchinsky and Wang [16] showed the following result.

Theorem A. *Let $\Omega \in Lip_\alpha(S^{n-1})$, $0 < \alpha \leq 1$. If $w \in A_p$ (Muckenhoupt weight class), $1 < p < \infty$, then there exists a constant C independent of f such that*

$$\|\mu_\Omega(f)\|_{L_w^p} \leq C\|f\|_{L_w^p}.$$

Assume that Ω satisfies the same conditions as above, in [2] and [7], the authors proved the H_w^p - L_w^p boundedness of Marcinkiewicz integrals provided that $\frac{n}{n+\beta} < p < 1$ and $w \in A_{p(1+\beta/n)}$, where $\beta = \min\{\alpha, 1/2\}$. The main purpose of this paper is to discuss the weak type estimate of μ_Ω on the weighted Hardy spaces $H_w^p(\mathbb{R}^n)$ when $p = \frac{n}{n+\alpha}$ and $w \in A_1$. In the meantime, the corresponding weak type estimate of μ_Ω on the homogeneous weighted Herz-type Hardy spaces $HK_q^{\alpha,p}(w_1, w_2)$ is also given. We now state our main results as follows.

Theorem 1. *Let $0 < \alpha < 1$ and $\Omega \in Lip_\alpha(S^{n-1})$. If $p = \frac{n}{n+\alpha}$, $w \in A_1$, then there exists a constant $C > 0$ independent of f such that*

$$\|\mu_\Omega(f)\|_{WL_w^p} \leq C\|f\|_{H_w^p}.$$

Theorem 2. *Let $0 < \beta < 1$ and $\Omega \in Lip_\beta(S^{n-1})$. If $0 < p \leq 1$, $1 < q < \infty$, $\alpha = n(1 - 1/q) + \beta$, $w_1, w_2 \in A_1$, then there exists a constant C independent of f such that*

$$\|\mu_\Omega(f)\|_{WK_q^{\alpha,p}(w_1, w_2)} \leq C\|f\|_{HK_q^{\alpha,p}(w_1, w_2)}.$$

2 Notations and definitions

First, let's recall some standard definitions and notations. The classical A_p weight theory was first introduced by Muckenhoupt in the study of weighted L^p boundedness of Hardy-Littlewood maximal functions in [13]. Let w be a nonnegative, locally integrable function defined on $\mathbb{R}^n$, all cubes are assumed

to have their sides parallel to the coordinate axes. We say that $w \in A_p$, $1 < p < \infty$, if

$$\left(\frac{1}{|Q|} \int_Q w(x) dx \right) \left(\frac{1}{|Q|} \int_Q w(x)^{-\frac{1}{p-1}} dx \right)^{p-1} \leq C \quad \text{for every cube } Q \subseteq \mathbb{R}^n,$$

where C is a positive constant which is independent of the choice of Q .

For the case $p = 1$, $w \in A_1$, if

$$\frac{1}{|Q|} \int_Q w(x) dx \leq C \cdot \operatorname{ess\,inf}_{x \in Q} w(x) \quad \text{for every cube } Q \subseteq \mathbb{R}^n.$$

A weight function w is said to belong to the reverse Hölder class RH_r if there exist two constants $r > 1$ and $C > 0$ such that the following reverse Hölder inequality holds

$$\left(\frac{1}{|Q|} \int_Q w(x)^r dx \right)^{1/r} \leq C \left(\frac{1}{|Q|} \int_Q w(x) dx \right) \quad \text{for every cube } Q \subseteq \mathbb{R}^n.$$

It is well known that if $w \in A_p$ with $1 < p < \infty$, then $w \in A_r$ for all $r > p$, and $w \in A_q$ for some $1 < q < p$. We thus write $q_w \equiv \inf\{q > 1 : w \in A_q\}$ to denote the critical index of w . If $w \in A_p$ with $1 \leq p < \infty$, then there exists $r > 1$ such that $w \in RH_r$.

Given a cube Q and $\lambda > 0$, λQ denotes the cube with the same center as Q whose side length is λ times that of Q . $Q = Q(x_0, r_Q)$ denotes the cube centered at x_0 with side length r_Q . For a weight function w and a measurable set E , we set the weighted measure $w(E) = \int_E w(x) dx$.

We shall need the following lemmas.

Lemma B ([5]). *Let $w \in A_p$, $p \geq 1$. Then, for any cube Q , there exists an absolute constant $C > 0$ such that*

$$w(2Q) \leq C w(Q).$$

In general, for any $\lambda > 1$, we have

$$w(\lambda Q) \leq C \cdot \lambda^{np} w(Q),$$

where C does not depend on Q nor on λ .

Lemma C ([5,6]). *Let $w \in A_p \cap RH_r$, $p \geq 1$ and $r > 1$. Then there exist constants $C_1, C_2 > 0$ such that*

$$C_1 \left(\frac{|E|}{|Q|} \right)^p \leq \frac{w(E)}{w(Q)} \leq C_2 \left(\frac{|E|}{|Q|} \right)^{(r-1)/r}$$

for any measurable subset E of a cube Q .

It should be pointed out that the definition of A_p ($1 \leq p < \infty$) condition could have been given with balls B replacing the cubes Q and the conclusions of Lemmas B and C also hold.

Next we shall give the definitions of the weighted Hardy spaces $H_w^p(\mathbb{R}^n)$ and homogeneous weighted Herz-type Hardy spaces $\dot{K}_q^{\alpha,p}(w_1, w_2)$. Given a Muckenhoupt's weight function w on $\mathbb{R}^n$, for $0 < p < \infty$, we denote by $L_w^p(\mathbb{R}^n)$ the space of all functions satisfying

$$\|f\|_{L_w^p(\mathbb{R}^n)} = \left(\int_{\mathbb{R}^n} |f(x)|^p w(x) dx \right)^{1/p} < \infty.$$

We also denote by $WL_w^p(\mathbb{R}^n)$ the weak weighted L^p space which is formed by all functions satisfying

$$\|f\|_{WL_w^p(\mathbb{R}^n)} = \sup_{\lambda > 0} \lambda \cdot w(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})^{1/p} < \infty.$$

Let $\mathcal{S}(\mathbb{R}^n)$ be the class of Schwartz functions and let $\mathcal{S}'(\mathbb{R}^n)$ be its dual space. Suppose that φ is a function in $\mathcal{S}'(\mathbb{R}^n)$ satisfying $\int_{\mathbb{R}^n} \varphi(x) dx = 1$. Set

$$\varphi_t(x) = t^{-n} \varphi(x/t), \quad t > 0, x \in \mathbb{R}^n.$$

For $f \in \mathcal{S}'(\mathbb{R}^n)$, we will define the maximal function $M_\varphi f(x)$ by

$$M_\varphi f(x) = \sup_{t > 0} |f * \varphi_t(x)|.$$

Definition 1. Let $0 < p < \infty$ and w be a weight function on $\mathbb{R}^n$. Then the weighted Hardy space $H_w^p(\mathbb{R}^n)$ is defined by

$$H_w^p(\mathbb{R}^n) = \{f \in \mathcal{S}'(\mathbb{R}^n) : M_\varphi f \in L_w^p(\mathbb{R}^n)\}$$

and we define $\|f\|_{H_w^p} = \|M_\varphi f\|_{L_w^p}$.

Set $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ and $C_k = B_k \setminus B_{k-1}$ for $k \in \mathbb{Z}$. Denote $\chi_k = \chi_{C_k}$ for $k \in \mathbb{Z}$, $\tilde{\chi}_k = \chi_k$ if $k \in \mathbb{N}$ and $\tilde{\chi}_0 = \chi_{B_0}$, where χ_{C_k} is the characteristic function of C_k . Let $\alpha \in \mathbb{R}$, $0 < p, q < \infty$ and w_1, w_2 be two weight functions on $\mathbb{R}^n$. The homogeneous weighted Herz space $\dot{K}_q^{\alpha,p}(w_1, w_2)$ is defined by

$$\dot{K}_q^{\alpha,p}(w_1, w_2) = \{f \in L_{loc}^q(\mathbb{R}^n \setminus \{0\}, w_2) : \|f\|_{\dot{K}_q^{\alpha,p}(w_1, w_2)} < \infty\},$$

where

$$\|f\|_{\dot{K}_q^{\alpha,p}(w_1, w_2)} = \left(\sum_{k \in \mathbb{Z}} (w_1(B_k))^{\alpha p/n} \|f \chi_k\|_{L_{w_2}^q}^p \right)^{1/p}.$$

For $k \in \mathbb{Z}$ and $\lambda > 0$, we set $E_k(\lambda, f) = |\{x \in C_k : |f(x)| > \lambda\}|$. Let $\tilde{E}_k(\lambda, f) = E_k(\lambda, f)$ for $k \in \mathbb{N}$ and $E_0(\lambda, f) = |\{x \in B(0, 1) : |f(x)| > \lambda\}|$. A measurable function $f(x)$ on $\mathbb{R}^n$ is said to belong to the homogeneous weak weighted Herz space $W\dot{K}_q^{\alpha,p}(w_1, w_2)$ if

$$\|f\|_{W\dot{K}_q^{\alpha,p}(w_1, w_2)} = \sup_{\lambda > 0} \lambda \cdot \left(\sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} w_2(E_k(\lambda, f))^{p/q} \right)^{1/p} < \infty.$$

For $f \in \mathcal{S}'(\mathbb{R}^n)$, the grand maximal function of f is defined by

$$G(f)(x) = \sup_{\varphi \in \mathcal{A}_N} \sup_{|y-x| < t} |\varphi_t * f(y)|,$$

where $N > n + 1$, $\mathcal{A}_N = \{\varphi \in \mathcal{S}(\mathbb{R}^n) : \sup_{|\alpha|, |\beta| \leq N} |x^\alpha D^\beta \varphi(x)| \leq 1\}$.

Definition 2. Let $0 < \alpha < \infty$, $0 < p < \infty$, $1 < q < \infty$ and w_1, w_2 be two weight functions on $\mathbb{R}^n$. The homogeneous weighted Herz-type Hardy space $H\dot{K}_q^{\alpha,p}(w_1, w_2)$ associated with the space $\dot{K}_q^{\alpha,p}(w_1, w_2)$ is defined by

$$H\dot{K}_q^{\alpha,p}(w_1, w_2) = \{f \in \mathcal{S}'(\mathbb{R}^n) : G(f) \in \dot{K}_q^{\alpha,p}(w_1, w_2)\}$$

and we define $\|f\|_{H\dot{K}_q^{\alpha,p}(w_1, w_2)} = \|G(f)\|_{\dot{K}_q^{\alpha,p}(w_1, w_2)}$.

3 The atomic decomposition

In this section, we will give the atomic decomposition theorems for weighted Hardy spaces and homogeneous weighted Herz-type Hardy spaces. In [4], Garcia-Cuerva characterized weighted Hardy spaces in terms of atoms in the following way.

Definition 3. Let $0 < p \leq 1 \leq q \leq \infty$ and $p \neq q$ such that $w \in A_q$ with critical index q_w . Set $[\cdot]$ the greatest integer function. For $s \in \mathbb{Z}_+$ satisfying $s \geq [n(q_w/p - 1)]$, a real-valued function $a(x)$ is called (p, q, s) -atom centered at x_0 with respect to w (or w -(p, q, s)-atom centered at x_0) if the following conditions are satisfied:

- (a) $a \in L_w^q(\mathbb{R}^n)$ and is supported in a cube Q centered at x_0 ,
- (b) $\|a\|_{L_w^q} \leq w(Q)^{1/q-1/p}$,
- (c) $\int_{\mathbb{R}^n} a(x) x^\alpha dx = 0$ for every multi-index α with $|\alpha| \leq s$.

Theorem D. Let $0 < p \leq 1 \leq q \leq \infty$ and $p \neq q$ such that $w \in A_q$ with critical index q_w . For each $f \in H_w^p(\mathbb{R}^n)$, there exist a sequence $\{a_j\}$ of w -($p, q, [n(q_w/p - 1)]$)-atoms and a sequence $\{\lambda_j\}$ of real numbers with $\sum_j |\lambda_j|^p \leq C \|f\|_{H_w^p}^p$ such that $f = \sum_j \lambda_j a_j$ both in the sense of distributions and in the H_w^p norm.

In [10] and [11], Lu and Yang characterized homogeneous weighted Herz-type Hardy spaces in terms of atoms as follows.

Definition 4. Let $1 < q < \infty$, $n(1-1/q) \leq \alpha < \infty$ and $s \geq [\alpha + n(1/q-1)]$. A real-valued function $a(x)$ is called a central (α, q, s) -atom with respect to (w_1, w_2) (or a central $(\alpha, q, s; w_1, w_2)$ -atom), if it satisfies

- (a) $\text{supp } a \subseteq B(0, R) = \{x \in \mathbb{R}^n : |x| < R\}$,
- (b) $\|a\|_{L_{w_2}^q} \leq w_1(B(0, R))^{-\alpha/n}$,
- (c) $\int_{\mathbb{R}^n} a(x) x^\beta dx = 0$ for every multi-index β with $|\beta| \leq s$.

Theorem E. Let $w_1, w_2 \in A_1$, $0 < p < \infty$, $1 < q < \infty$ and $n(1-1/q) \leq \alpha < \infty$. Then we have that $f \in H\dot{K}_q^{\alpha, p}(w_1, w_2)$ if and only if

$$f(x) = \sum_{k \in \mathbb{Z}} \lambda_k a_k(x), \quad \text{in the sense of } \mathcal{S}'(\mathbb{R}^n),$$

where $\sum_{k \in \mathbb{Z}} |\lambda_k|^p < \infty$, each a_k is a central $(\alpha, q, s; w_1, w_2)$ -atom. Moreover,

$$\|f\|_{H\dot{K}_q^{\alpha, p}(w_1, w_2)} \approx \inf \left(\sum_{k \in \mathbb{Z}} |\lambda_k|^p \right)^{1/p},$$

where the infimum is taken over all the above decompositions of f .

For the properties and applications of the above two spaces, we refer the readers to the books [12] and [15] for further details. Throughout this article, we will use C to denote a positive constant, which is independent of the main parameters and not necessarily the same at each occurrence. By $A \sim B$, we mean that there exists a constant $C > 1$ such that $\frac{1}{C} \leq \frac{A}{B} \leq C$. Moreover, we will denote the conjugate exponent of $q > 1$ by $q' = q/(q-1)$.

4 Proof of Theorem 1

In order to prove our main result, we shall need the following superposition principle on the weighted weak type estimates.

Theorem 4.1. Let $w \in A_1$ and $0 < p < 1$. If a sequence of measurable functions $\{f_j\}$ satisfy

$$\|f_j\|_{WL_w^p} \leq 1 \quad \text{for all } j \in \mathbb{Z}$$

and

$$\sum_{j \in \mathbb{Z}} |\lambda_j|^p \leq 1,$$

then we have

$$\left\| \sum_{j \in \mathbb{Z}} \lambda_j f_j \right\|_{WL_w^p} \leq \left(\frac{2-p}{1-p} \right)^{1/p}.$$

Proof. The proof of this lemma is similar to the corresponding result for the unweighted case. See [8, page 123]. $\square$

We are now in a position to give the proof of Theorem 1.

Proof. We note that when $w \in A_1$ and $p = n/(n + \alpha)$, then $[n(q_w/p - 1)] = [\alpha] = 0$. By Lemma 4.1 and Theorem D, it suffices to show that for any w -($p, q, 0$)-atom a , there exists a constant $C > 0$ independent of a such that $\|\mu_\Omega(a)\|_{WL_w^p} \leq C$.

Let a be a w -($p, q, 0$)-atom with $\text{supp } a \subseteq Q = Q(x_0, r_Q)$, $1 < q < \infty$ and let $Q^* = 2\sqrt{n}Q$. For any given $\lambda > 0$, we can write

$$\begin{aligned} & \lambda^p \cdot w(\{x \in \mathbb{R}^n : |\mu_\Omega(a)(x)| > \lambda\}) \\ & \leq \lambda^p \cdot w(\{x \in Q^* : |\mu_\Omega(a)(x)| > \lambda\}) + \lambda^p \cdot w(\{x \in (Q^*)^c : |\mu_\Omega(a)(x)| > \lambda\}) \\ & = I_1 + I_2. \end{aligned}$$

Since $w \in A_1$, then $w \in A_q$ for $1 < q < \infty$. Applying Chebyshev's inequality, Hölder's inequality, Lemma B and Theorem A, we thus have

$$\begin{aligned} I_1 & \leq \int_{Q^*} |\mu_\Omega(a)(x)|^p w(x) dx \\ & \leq \left(\int_{Q^*} |\mu_\Omega(a)(x)|^q w(x) dx \right)^{p/q} \left(\int_{Q^*} w(x) dx \right)^{1-p/q} \\ & \leq \|\mu_\Omega(a)\|_{L_w^q}^p w(Q)^{1-p/q} \\ & \leq C \cdot \|a\|_{L_w^q}^p w(Q)^{1-p/q} \\ & \leq C. \end{aligned}$$

We now turn to estimate I_2 . If we set $\varphi(x) = \Omega(x)|x|^{-n+1}\chi_{\{|x| \leq 1\}}(x)$, then

$$\mu_\Omega(f)(x) = \left(\int_0^\infty |\varphi_t * f(x)|^2 \frac{dt}{t} \right)^{1/2}.$$

By the vanishing moment condition of atom a , we have

$$\begin{aligned}
|\varphi_t * a(x)| &= \frac{1}{t} \cdot \left| \int_Q \left(\frac{\Omega(x-y)}{|x-y|^{n-1}} - \frac{\Omega(x-x_0)}{|x-x_0|^{n-1}} \right) a(y) dy \right| \\
&\leq C \cdot \frac{1}{t} \int_Q \left| \frac{1}{|x-y|^{n-1}} - \frac{1}{|x-x_0|^{n-1}} \right| |a(y)| dy \\
&\quad + \frac{1}{t} \int_Q \frac{|\Omega(x-y) - \Omega(x-x_0)|}{|x-x_0|^{n-1}} |a(y)| dy \\
&= \text{I} + \text{II}.
\end{aligned}$$

Observe that when $y \in Q$, $x \in (Q^*)^c$, then $|x-y| \sim |x-x_0|$. This together with the mean value theorem gives

$$\text{I} \leq C \cdot \frac{rQ}{t|x-x_0|^n} \int_Q |a(y)| dy. \quad (1)$$

On the other hand, since $\Omega \in Lip_\alpha(S^{n-1})$, $0 < \alpha < 1$, then we can get

$$\begin{aligned}
\text{II} &\leq C \cdot \frac{1}{t|x-x_0|^{n-1}} \int_Q \left| \frac{x-y}{|x-y|} - \frac{x-x_0}{|x-x_0|} \right|^\alpha |a(y)| dy \\
&\leq C \cdot \frac{1}{t|x-x_0|^{n-1}} \int_Q \left(\frac{|y-x_0|}{|x-x_0|} \right)^\alpha |a(y)| dy \\
&\leq C \cdot \frac{(rQ)^\alpha}{t|x-x_0|^{n-1+\alpha}} \int_Q |a(y)| dy.
\end{aligned} \quad (2)$$

By using Hölder's inequality and the A_q condition, we thus obtain

$$\begin{aligned}
\int_Q |a(y)| dy &\leq \left(\int_Q |a(y)|^q w(y) dy \right)^{1/q} \left(\int_Q w(y)^{-q'/q} dy \right)^{1/q'} \\
&\leq C \cdot \|a\|_{L_w^q} \left(\frac{|Q|^q}{w(Q)} \right)^{1/q} \\
&\leq C \cdot \frac{|Q|}{w(Q)^{1/p}}.
\end{aligned} \quad (3)$$

We also observe that $\text{supp } \varphi \subseteq \{x \in \mathbb{R}^n : |x| \leq 1\}$, then for any $y \in Q$, $x \in (Q^*)^c$, we have $t \geq |x-y| \geq |x-x_0| - |y-x_0| \geq \frac{|x-x_0|}{2}$. Substituting

the above inequality (3) into (1) and (2), we can deduce

$$\begin{aligned}
|\mu_\Omega(a)(x)|^2 &\leq C \left(\frac{r_Q^{n+1}}{|x-x_0|^n w(Q)^{1/p}} + \frac{r_Q^{n+\alpha}}{|x-x_0|^{n-1+\alpha} w(Q)^{1/p}} \right)^2 \left(\int_{\frac{|x-x_0|}{2}}^\infty \frac{dt}{t^3} \right) \\
&\leq C \left(\frac{r_Q^{n+1}}{|x-x_0|^{n+1} w(Q)^{1/p}} + \frac{r_Q^{n+\alpha}}{|x-x_0|^{n+\alpha} w(Q)^{1/p}} \right)^2 \\
&\leq C \left(\frac{1}{w(Q)^{1/p}} \right)^2.
\end{aligned}$$

Set $Q_0^* = Q$, $Q_1^* = Q^*$ and $Q_k^* = (Q_{k-1}^*)^*$, $k = 2, 3, \dots$. Following along the same lines as above, we can also show that for any $x \in (Q_k^*)^c$, then

$$|\mu_\Omega(a)(x)| \leq C \cdot \frac{1}{w(Q_{k-1}^*)^{1/p}} \quad k = 1, 2, \dots$$

We shall consider the following two cases:

If $\{x \in (Q^*)^c : |\mu_\Omega(a)(x)| > \lambda\} = \emptyset$, then the inequality

$$I_2 \leq C$$

holds trivially.

If $\{x \in (Q^*)^c : |\mu_\Omega(a)(x)| > \lambda\} \neq \emptyset$, then for $p = n/(n+\alpha)$, it is easy to check that

$$\lim_{k \rightarrow \infty} \frac{1}{w(Q_k^*)^{1/p}} = 0.$$

Consequently, for any fixed $\lambda > 0$, we are able to find a maximal positive integer N such that

$$\lambda < C \cdot \frac{1}{w(Q_N^*)^{1/p}}.$$

Therefore

$$\begin{aligned}
I_2 &\leq \lambda^p \cdot \sum_{k=1}^N w(\{x \in Q_{k+1}^* \setminus Q_k^* : |\mu_\Omega(a)(x)| > \lambda\}) \\
&\leq C \cdot \frac{1}{w(Q_N^*)} \sum_{k=1}^N w(Q_{k+1}^*) \\
&\leq C.
\end{aligned}$$

Combining the above estimates for I_1 , I_2 and taking the supremum over all $\lambda > 0$, we complete the proof of Theorem 1. $\square$

5 Proof of Theorem 2

Proof. We note that our assumption $\alpha = n(1 - 1/q) + \beta$ implies that $s = [\alpha + n(1/q - 1)] = [\beta] = 0$. For every $f \in H\dot{K}_q^{\alpha,p}(w_1, w_2)$, then by Theorem E, we have the decomposition $f = \sum_{j \in \mathbb{Z}} \lambda_j a_j$, where $\sum_{j \in \mathbb{Z}} |\lambda_j|^p < \infty$ and each a_j is a central $(\alpha, q, 0; w_1, w_2)$ -atom. Without loss of generality, we may assume that $\text{supp } a_j \subseteq B(0, R_j)$ and $R_j = 2^j$. For any given $\sigma > 0$, we write

$$\begin{aligned} & \sigma^p \cdot \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} w_2(\{x \in C_k : |\mu_\Omega(f)(x)| > \sigma\})^{p/q} \\ & \leq \sigma^p \cdot \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} w_2(\{x \in C_k : \sum_{j=k-1}^{\infty} |\lambda_j| |\mu_\Omega(a_j)(x)| > \sigma/2\})^{p/q} \\ & \quad + \sigma^p \cdot \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} w_2(\{x \in C_k : \sum_{j=-\infty}^{k-2} |\lambda_j| |\mu_\Omega(a_j)(x)| > \sigma/2\})^{p/q} \\ & = J_1 + J_2. \end{aligned}$$

Since $w_2 \in A_1$, then $w_2 \in A_q$ for any $1 < q < \infty$. Note that $0 < p \leq 1$, then by using Chebyshev's inequality and Theorem A, we can get

$$\begin{aligned} J_1 & \leq 2^p \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} \left(\sum_{j=k-1}^{\infty} |\lambda_j| \|\mu_\Omega(a_j) \chi_k\|_{L_{w_2}^q} \right)^p \\ & \leq 2^p \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} \left(\sum_{j=k-1}^{\infty} |\lambda_j|^p \|\mu_\Omega(a_j)\|_{L_{w_2}^q}^p \right) \\ & \leq C \sum_{k \in \mathbb{Z}} w_1(B_k)^{\alpha p/n} \left(\sum_{j=k-1}^{\infty} |\lambda_j|^p \|a_j\|_{L_{w_2}^q}^p \right). \end{aligned}$$

Changing the order of summation yields

$$J_1 \leq C \sum_{j \in \mathbb{Z}} |\lambda_j|^p \left(\sum_{k=-\infty}^{j+1} w_1(B_k)^{\alpha p/n} w_1(B_j)^{-\alpha p/n} \right).$$

When $k \leq j+1$, then $B_k \subseteq B_{j+1}$. Since $w_1 \in A_1$, then we know $w \in RH_r$ for some $r > 1$. It follows directly from Lemma C that

$$w_1(B_k) \leq C \cdot w_1(B_{j+1}) |B_k|^\delta |B_{j+1}|^{-\delta}, \quad (4)$$

where $\delta = (r-1)/r > 0$. By Lemma B and the above inequality (4), we get

$$\begin{aligned}
& \sum_{k=-\infty}^{j+1} w_1(B_k)^{\alpha p/n} w_1(B_j)^{-\alpha p/n} \\
& \leq C \sum_{k=-\infty}^{j+1} \left(\frac{w_1(B_{j+1})}{w_1(B_j)} \right)^{\alpha p/n} \left(\frac{|B_k|}{|B_{j+1}|} \right)^{\alpha \delta p/n} \\
& \leq C \sum_{k=-\infty}^{j+1} 2^{(k-j-1)\alpha \delta p} \\
& \leq C \sum_{k=0}^{\infty} 2^{-k\alpha \delta p},
\end{aligned}$$

where the last series is convergent since $\alpha \delta p > 0$. Furthermore, it is bounded by a constant which is independent of $j \in \mathbb{Z}$. Hence

$$J_1 \leq C \sum_{j \in \mathbb{Z}} |\lambda_j|^p \leq C \|f\|_{H\dot{K}_q^{\alpha,p}(w_1,w_2)}^p.$$

We turn to deal with J_2 . As in the proof of Theorem 1, we can also write

$$\begin{aligned}
|\varphi_t * a_j(x)| &= \frac{1}{t} \cdot \left| \int_{B_j} \left(\frac{\Omega(x-y)}{|x-y|^{n-1}} - \frac{\Omega(x)}{|x|^{n-1}} \right) a_j(y) dy \right| \\
&\leq C \cdot \frac{1}{t} \int_{B_j} \left| \frac{1}{|x-y|^{n-1}} - \frac{1}{|x|^{n-1}} \right| |a_j(y)| dy \\
&\quad + \frac{1}{t} \int_{B_j} \frac{|\Omega(x-y) - \Omega(x)|}{|x|^{n-1}} |a_j(y)| dy \\
&= \text{III} + \text{IV}.
\end{aligned}$$

Observe that when $j \leq k-2$, then for any $y \in B_j$ and $x \in C_k = B_k \setminus B_{k-1}$, we have $|x| \geq 2|y|$, which implies $|x-y| \sim |x|$. We also observe that $\text{supp } \varphi \subseteq \{x \in \mathbb{R}^n : |x| \leq 1\}$, then we can get $t \geq |x-y| \geq \frac{|x|}{2}$. Hence, by using the same arguments as that of Theorem 1, we obtain

$$\text{III} \leq C \cdot \frac{R_j}{t|x|^n} \int_{B_j} |a_j(y)| dy \quad (5)$$

and

$$\text{IV} \leq C \cdot \frac{(R_j)^\beta}{t|x|^{n-1+\beta}} \int_{B_j} |a_j(y)| dy. \quad (6)$$

Similarly, it follows from Hölder's inequality and the A_q condition that

$$\begin{aligned} \int_{B_j} |a_j(y)| dy &\leq \left(\int_{B_j} |a_j(y)|^q w_2(y) dy \right)^{1/q} \left(\int_{B_j} w_2(y)^{-q'/q} dy \right)^{1/q'} \\ &\leq C \cdot |B_j| w_1(B_j)^{-\alpha/n} w_2(B_j)^{-1/q}. \end{aligned} \quad (7)$$

Substituting the above inequality (7) into (5) and (6), we can deduce

$$\begin{aligned} &|\mu_\Omega(a_j)(x)|^2 \\ &\leq C \left(\frac{2^{j(n+1)}}{|x|^n w_1(B_j)^{\alpha/n} w_2(B_j)^{1/q}} + \frac{2^{j(n+\beta)}}{|x|^{n-1+\beta} w_1(B_j)^{\alpha/n} w_2(B_j)^{1/q}} \right)^2 \left(\int_{\frac{|x|}{2}}^\infty \frac{dt}{t^3} \right) \\ &\leq C \left(\frac{2^{j(n+1)}}{|x|^{n+1} w_1(B_j)^{\alpha/n} w_2(B_j)^{1/q}} + \frac{2^{j(n+\beta)}}{|x|^{n+\beta} w_1(B_j)^{\alpha/n} w_2(B_j)^{1/q}} \right)^2. \end{aligned} \quad (8)$$

Since $B_j \subseteq B_{k-2}$, then by using Lemma C, we get

$$w_i(B_j) \geq C \cdot w_i(B_{k-2}) |B_j| |B_{k-2}|^{-1} \quad \text{for } i = 1 \text{ or } 2.$$

From our assumption $\alpha = n(1 - 1/q) + \beta$ and (8), it follows that

$$\begin{aligned} |\mu_\Omega(a_j)(x)| &\leq C \cdot \left(\frac{2^j}{2^{k-2}} \right)^{n+\beta-\alpha-n/q} \frac{1}{w_1(B_{k-2})^{\alpha/n} w_2(B_{k-2})^{1/q}} \\ &\leq C \cdot \frac{1}{w_1(B_{k-2})^{\alpha/n} w_2(B_{k-2})^{1/q}}. \end{aligned} \quad (9)$$

We now set $A_k = w_1(B_{k-2})^{-\alpha/n} w_2(B_{k-2})^{-1/q}$. Once again, let us consider the following two cases:

If $\{x \in C_k : \sum_{j=-\infty}^{k-2} |\lambda_j| |\mu_\Omega(a_j)(x)| > \sigma/2\} = \emptyset$, then the inequality

$$J_2 \leq C \|f\|_{H\dot{K}_q^{\alpha,p}(w_1,w_2)}^p$$

holds trivially.

If $\{x \in C_k : \sum_{j=-\infty}^{k-2} |\lambda_j| |\mu_\Omega(a_j)(x)| > \sigma/2\} \neq \emptyset$, then by the inequality (9), we have

$$\begin{aligned} \sigma &< C \cdot A_k \left(\sum_{j \in \mathbb{Z}} |\lambda_j| \right) \\ &\leq C \cdot A_k \left(\sum_{j \in \mathbb{Z}} |\lambda_j|^p \right)^{1/p} \\ &\leq C \cdot A_k \|f\|_{H\dot{K}_q^{\alpha,p}(w_1,w_2)}. \end{aligned}$$

In addition, it is easy to verify that $\lim_{k \rightarrow \infty} A_k = 0$. Then for any given $\sigma > 0$, we are able to find a maximal positive integer k_σ such that

$$\sigma < C \cdot A_{k_\sigma} \|f\|_{\dot{H}K_q^{\alpha,p}(w_1, w_2)}.$$

From the above discussion, we have that $B_{k-2} \subseteq B_{k_\sigma-2}$. As (4), by using Lemma C again, we obtain

$$\frac{w_i(B_{k-2})}{w_i(B_{k_\sigma-2})} \leq C \left(\frac{|B_{k-2}|}{|B_{k_\sigma-2}|} \right)^\delta \quad \text{for } i = 1 \text{ or } 2.$$

Furthermore, it follows immediately from Lemma B that

$$\frac{w_i(B_k)}{w_i(B_{k_\sigma-2})} \leq C \left(\frac{|B_{k-2}|}{|B_{k_\sigma-2}|} \right)^\delta \quad \text{for } i = 1 \text{ or } 2.$$

Therefore

$$\begin{aligned} J_2 &\leq \sigma^p \cdot \sum_{k=-\infty}^{k_\sigma} w_1(B_k)^{\alpha p/n} w_2(B_k)^{p/q} \\ &\leq C \|f\|_{\dot{H}K_q^{\alpha,p}(w_1, w_2)}^p \sum_{k=-\infty}^{k_\sigma} \left(\frac{w_1(B_k)}{w_1(B_{k_\sigma-2})} \right)^{\alpha p/n} \left(\frac{w_2(B_k)}{w_2(B_{k_\sigma-2})} \right)^{p/q} \\ &\leq C \|f\|_{\dot{H}K_q^{\alpha,p}(w_1, w_2)}^p \sum_{k=-\infty}^{k_\sigma} \frac{1}{2^{(k_\sigma-k)n\delta}} \\ &\leq C \|f\|_{\dot{H}K_q^{\alpha,p}(w_1, w_2)}^p. \end{aligned}$$

Finally, by combining the above estimates for J_1 , J_2 and taking the supremum over all $\sigma > 0$, we conclude the proof of Theorem 2. $\square$

References

- [1] Y. Ding, D. Fan, Y. Pan, Weighted boundedness for a class of rough Marcinkiewicz integrals, *Indiana Univ. Math. J.*, **48**(1999), 1037–1055.
- [2] Y. Ding, M. Y. Lee and C. C. Lin, Marcinkiewicz integral on weighted Hardy spaces, *Arch. Math.*, **80**(2003), 620–629.
- [3] Y. Ding, S. Lu, K. Yabuta, On commutators of Marcinkiewicz integrals with rough kernel, *J. Math. Anal. Appl.*, **275**(2002), 60–68.

- [4] J. Garcia-Cuerva, Weighted H^p spaces, *Dissertations Math*, **162**(1979), 1–63.
- [5] J. Garcia-Cuerva and J. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland, Amsterdam, 1985.
- [6] R. F. Gundy and R. L. Wheeden, Weighted integral inequalities for nontangential maximal function, Lusin area integral, and Walsh-Paley series, *Studia Math*, **49**(1974), 107–124.
- [7] C. C. Lin and Y. C. Lin, H_w^p - L_w^p boundedness of Marcinkiewicz integral, *Integr. equ. oper. theory*, **58**(2007), 87–98.
- [8] S. Lu, *Four Lectures on Real H^p Spaces*, World Scientific Publishing, River Edge, N.J., 1995.
- [9] S. Lu, Marcinkiewicz integral with rough kernels, *Front. Math. China*, **3**(1)(2008), 1–14.
- [10] S. Lu and D. Yang, The decomposition of the weighted Herz spaces and its applications, *Sci. China(Ser. A)*, **38**(1995), 147–158.
- [11] S. Lu and D. Yang, The weighted Herz-type Hardy spaces and its applications, *Sci. China(Ser. A)*, **38**(1995), 662–673.
- [12] S. Lu, D. Yang and G. Hu, *Herz Type Spaces and Their Applications*, Science Press, Beijing, 2008.
- [13] B. Muckenhoupt, Weighted norm inequalities for the Hardy maximal function, *Trans. Amer. Math. Soc*, **165**(1972), 207–226.
- [14] E. M. Stein, On the functions of Littlewood-Paley, Lusin and Marcinkiewicz, *Trans. Amer. Math. Soc*, **88**(1958), 430–466.
- [15] J. O. Stömberg and A. Torchinsky, *Weighted Hardy spaces*, *Lecture Notes in Math*, Vol 1381, Springer-Verlag, 1989.
- [16] A. Torchinsky and S. Wang, A note on the Marcinkiewicz integral, *Colloq. Math*, **60/61**(1990), 235–243.